

and Gredos, we only found statistically significant differences in LIA ($p < 0.01$), SDI ($p < 0.01$) and M_1-M_3 ($p < 0.01$). Between the Galicia and Guadarrama samples, the most marked divergences were in α ($p < 0.05$), LIA ($p < 0.01$) and SDI ($p < 0.01$). Between Gredos and Guadarrama, there was an even greater degree of biometric similarity, only α showing significant differences ($p < 0.01$).

In view of the mean values obtained for all these populations (see Table 3), the specimens from Galicia generally evidence the largest mandibular dimensions, although these differences lack statistical significance (observe values of β , LIA, LM, SDI, LFT, M_1-M_3 , LC and HC). Due to their biometrical similarity, the relationship between the sizes of the Gredos and Guadarrama samples has not been completely defined.

To determine the degree of intersample affinity in each parameter, mean values were processed by the Student-Newman-Keuls test. In the graphic representation of results (Table 4), the populations with mean values not significantly heterogeneous are underlined (SOKAL and ROHLF 1979). The resulting diagrams evidence the close relationship among the populations analysed. The only variable to statistically separate the Guadarrama population was α , which was smaller in this sample. The specimens from Galicia differ from Gredos and Guadarrama in LIA, SDI and M_1-M_3 , the mean values of these parameters being higher in the Galicia samples.

On the basis of these findings, although there was no evidence of intraspecific mandibular biometric differences in *Sorex granarius*, there was a general tendency to smaller size from north to south. It is interesting to point out that in the Iberian Peninsula, *Crocidura russula* and *Crocidura suaveolens* have been found to present a clinal variation consisting in progressive reduction of craniometric values to the south (REY and LANDIN 1973; REY and REY 1974; SANS-COMA et al. 1987).

Acknowledgements

We are grateful to M. NAVARRO and Dr. V. PEREZ-MELLADO, who contributed 7 specimens and six new localities, as to Dr B. THOMAS, who translated this manuscript to English, and I. MAIER for translating to German.

Zusammenfassung

Verbreitung und Biometrie von Sorex granarius (Miller, 1910) (Soricinae: Insectivora)

Die Verbreitung von *Sorex granarius* ist auf die Pyrenäen-Halbinsel beschränkt. Hier ist diese Art in Gallizien, der nördlichen Hälfte von Portugal und im Zentralgebirge anzutreffen. *Sorex granarius* bewohnt vorwiegend Waldgebiete. Die vertikale Verbreitung erstreckt sich von der Meereshöhe bis zur Höhe von 2000 m. Im Zentralgebirge ist dieses Intervall kleiner (500–2000 m). Die Biometrie der Schädel von untersuchten Individuen ist sehr gleichartig, obwohl sich eine Tendenz zur Größenabnahme von Norden nach Süden beobachten lässt. Die größten Schädel wurden in Gallizien gefunden.

Literature

- ALMAÇA, C. (1968): La faune mammalogique du Portugal dans la checklist of Palearctic and Indian Mammals, 1951, par Ellerman et Morrison-Scott. Arq. Mus. Boc., 2 série, 12.
- ARENAS, A. (1983): Atlas de Micromamíferos de Segovia. Un modelo de análisis zoogeográfico de Insectívoros y Roedores mediante egagrópilas de lechuza. Tesina, Univ. Madrid.
- CABRERA, A. (1914): Fauna Ibérica. Mamíferos. Mus. Nac. Ciencias Nat.: Madrid.
- CAMPOS, F. (1977): Régimen alimenticio de *Tyto alba* en las provincias de Salamanca y Zamora. Ardeola 24, 105–119.
- CATZEFILIS, F. (1984): Systematique biochimique. Taxonomie et phylogénie des musaraignes d'Europe (Soricidae, Mammalia). Thèse, Université Lausanne.
- CATZEFILIS, F.; GRAF, J.-D.; HAUSSER, J.; VOGEL, P. (1982): Comparaison biochimique des Musaraignes du genre *Sorex* en Europe occidentale (Soricidae, Mammalia). Z. zool. Syst. Evolut.-forsch. 20, 223–233.
- DIXON, W. J. (1983): BMDP Statistical Software. Berkeley: University of California Press.

- GARZON-HEYDT, J.; Castroviejo, S.; Castroviejo, J. (1971): Notas preliminares sobre la distribución de algunos micromamíferos en el norte de España. *SäugetierkdL. Mitt.* 19, 217–222.
- HAUSSER, J. (1976): Contribution à l'étude des Musaraignes du genre *Sorex* (Cytotaxonomie, Morphologie, Répartition). Thèse, Université Lausanne.
- (1984): Genetic drift and selection: Their respective weights in the morphological and genetic differentiation of four species of shrews in Southern Europe (Insectivora, Soricidae). *Z. zool. Syst. Evolut.-forsch.* 22, 302–320.
- HAUSSER, J.; CATZEFLIS, F.; MEYLAN, A.; VOGEL, P. (1985): Speciation in the *Sorex araneus* complex (Mammalia: Insectivora). *Acta Zool. Fennica* 170, 125–130.
- HAUSSER, J.; GRAF, J.-D.; MEYLAN, A. (1975): Données nouvelles sur les *Sorex* d'Espagne et des Pyrénées (Mammalia, Insectivora). *Bull. Soc. Vaud. Sc. Nat.* 348, 241–252.
- HAUSSER, J.; JAMMOT, D. (1974): Etude biometrique des mâchoires chez les *Sorex* du groupe *araneus* en Europe continentale (Mammalia, Insectivora). *Bull. Soc. Vaud. Sc. Nat.* 348, 241–252.
- HEIM DE BALSAC, H.; DE BEAUFORT, F. (1969): Contribution à l'étude des micromammifères du Nord-Ouest de l'Espagne (Santander, Asturies, Galice, Leon). *Mammalia* 33, 367–380.
- JAMMOT, D. (1973): Mise au point d'un "bloc comparateur optique" pour les nécessités d'une ostéométrie fine. *Mammalia* 37, 122–125.
- LOPEZ-FUSTER, M. J. (1983): Sobre los géneros *Sorex* Linnaeus, 1758, *Suncus* Ehrenberg, 1833 y *Crocidura* Wagler, 1832 (Insectivora, Soricidae) en el Nordeste de la Península Ibérica. Tesis Doctoral, Universidad Barcelona.
- MADUREIRA, M. L.; MAGALHÃES, C. M. (1980): Small mammals of Portugal. *Arq. Mus. Bocage*, 2 ser., 7, 179–214.
- MADUREIRA, M. L.; RAMALINHO, M. G. (1981): Notas sobre a distribuição, diagnose e ecologia dos Insectívora e Rodentia portugueses. *Arq. Mus. Bocage*, ser. A, 1, 165–263.
- MALEC, F.; STORCH, G. (1964): Einige Kleinsäuger (Mammalia: Insectivora, Rodentia) aus Nordspanien. *Z. Säugetierkunde* 29, 220–230.
- MEYLAN, A.; HAUSSER, J. (1973): Les chromosomes des *Sorex* du groupe *araneus-arcticus* (Mammalia, Insectivora). *Z. Säugetierkunde* 38, 143–158.
- MILLER, G. S. (1910): Description of six new European Mammals. *Ann. Mag. Nat. Hist.*, Ser. 8, 6, 458–461.
- (1912): Catalogue of the Mammals of Western Europe. London: British Museum (Nat. Hist.).
- NIETHAMMER, J. (1956): Insektenfresser und Nager Spaniens. *Bonn. Zool. Beitr.* 7, 249–295.
- (1970): Über Kleinsäuger aus Portugal. *Bonn. Zool. Beitr.* 21, 89–118.
- NORES, C. (1979): Nuevas aportaciones al conocimiento de la subfamilia Soricinae (Mammalia, Insectivora) en los distritos Cantábrico y Lusitano. Tesis Licenciatura, Universidad Oviedo.
- RAMALINHO, M. G. (1981): Portuguese Soricinae. A review of recent data (in press).
- REY, J. M.; LANDIN, A. (1973): Sobre la presencia de *Crocidura suaveolens* en el sur de Andalucía (Mammalia, Insectivora). *Bol. R. Soc. Esp. Hist. Nat. (Biol.)* 71, 9–16.
- REY, J. C.; REY, J. M. (1974): Nota preliminar sobre las musarañas del género *Crocidura* Wagler, 1832 en las islas Baleares. *Bol. Est. Centr. Ecol.* 3, 79–85.
- RIVAS-MARTINEZ, S. (1981): Les étages bioclimatiques de la végétation de la Péninsule Ibérique. *Anal. Jardín Botánico de Madrid* 37, 251–268.
- (1983): Pisos bioclimáticos de España. *Lazaroa* 5, 33–43.
- SANS-COMA, V. (1979): Beitrag zur Kenntnis der Waldspitzmaus, *Sorex araneus* Linné, 1758, in Katalonien, Spanien. *SäugetierkdL. Mitt.* 27, 96–106.
- SANS-COMA, V.; LOPEZ-FUSTER, M. J.; VARGAS, J. M. (1983): Sobre la musaraña común, *Crocidura russula* (Hermann, 1780) en el sur de la Península Ibérica. In: Mamíferos y Helmintos. Ed. by M. BACH. Barcelona (in press).
- SOKAL, R. R.; ROHLF, F. J. (1979): Biometria. Barcelona: Ed. Blume.

Authors' addresses: J. GIBERT and ROSA GARCÍA-PEREÀ, Unidad de Zoología Aplicada, El Encin, Apdo. 127, Alcalá de Henares, Madrid, Spain; M. J. LÓPEZ-FUSTER and J. VENTURA, Departamento de Vertebrados, Facultad de Biología, Av. Diagonal 657, E-08028 Barcelona, Spain

Notes on some bats from northern Thailand, with comments on the subgeneric status of *Myotis altarium*

B. R. BLOOD and D. A. McFARLANE

Section of Birds and Mammals, Los Angeles County Museum of Natural History, Los Angeles

Receipt of Ms. 06. 04. 1987

Abstract

Reported is the occurrence of four species of bats from extreme northern Thailand collected by members of the National Speleological Society, USA, Thailand Karst Hydrologic Expedition. Single specimens of *Aselliscus stoliczkanus*, *Ia io*, *Miniopterus macrodons*, and *Myotis altarium* were collected. This is the first record of *M. altarium* outside of China and only the second record for *I. io*. Although known from the area, our specimen of *A. stoliczkanus* does not match published color patterns of this species from Thailand. We provide the first published photograph of the skull of *M. altarium* and an expanded description of this species because previously published descriptions are inadequate to place this species at the subgeneric level. We compared *M. altarium* directly to members of three other subgenera of *Myotis* and conclude it belongs in the subgenus *Myotis*.

Introduction

The Thailand Karst Hydrologic Survey, fielded by members of the National Speleological Society between December 1983 and February 1984, provided several specimens of cave-dwelling bats from the limestone mountains of extreme northern Thailand. We report here three species of Vespertilionidae plus one species of Hipposideridae. We found published descriptions of one of the species collected, *Myotis altarium*, inadequate to properly place it at the subgeneric level. Therefore, we present an expanded description of *M. altarium*, as well as, the first photographs of its skull. Specimens have been deposited in the mammal collection of the Natural History Museum of Los Angeles County (LACM). The rhinolophids collected during this expedition are reported separately (McFARLANE and BLOOD 1986).

Results and discussion

Systematic accounts

Aselliscus stoliczkanus Dobson, 1871 (Hipposideridae)

Specimen collected; LACM 70301; collected by R. HEMPERLY, 24 Dec. 1983; male, in alcohol with skull extracted. Northern Thailand; Chang Dao Mountain, Chang Dao Cave. 19°23'N, 98°54'E. Some selected measurements (in mm) are head and body length, 43; tail length, 38; forearm length, 41; greatest length of skull, 18; hind foot length, 5; ear length, 8. LEKAGUL and MCNEELEY (1977) report this species to be uncommon, but wide spread. They further report that this species occurs in two color phases, a brown-gray phase and a yellow-red phase. The LACM specimen has long dorsal fur (7 mm) which is a creamy white at its base with rich dark brown tips. The ventral fur color is a silvery brown.

Miniopterus macrodens Maeda, 1982 (Vespertilionidae)

Specimen collected: LACM 70323; collected by R. HEMPERLY 12 Jan. 1984; male, in alcohol with skull extracted. Northern Thailand, Aung Kang Region, Big House Cave; 19°50'N, 99°10'E. Selected measurements (in mm) are as follows; total length, 107; forearm length, 50.5; greatest length of skull, 16.4; ear length 16; tragus length, 4.5; length of third metacarpal, 47.5; length of first phalanx of digit 3, 11.9; length of second phalanx of digit 3, 34.5. Our specimen is all black in color including the flight membranes. This species is part of the *M. maginater* group (R. PETERSON, pers. comm.)

Ia io Thomas, 1902 (Vespertilionidae)

Specimen collected: LACM 70321; collected by R. HEMPERLY 11 Jan. 1984; in alcohol. Northern Thailand, Aung Kang Region, Big House Cave. 19°50'N, 99°10'E. Some representative measurements (in mm) include head and body length, 90.6; tail length, 71.5; forearm length, 75.3; ear length, 26; tragus length, 8.5; length of metacarpal III, 68.3; length of third phalanx of digit III, 11.9. This specimen was collected by hand near the entrance of Big House Cave. The field notes indicate that it was not in the immediate company of other bats. However, three other species were collected in this same cave: *Rhinolophus malayanus*, *R. robinsoni*, and *M. altarium* (MFARLANE and BLOOD 1986). Only one other specimen (Museum of Comparative Zoology 3549) is known from northern Thailand, near Chang Dao (ALLEN and COOLIDGE 1940).

Myotis altarium Thomas, 1911 (Vespertilionidae)

Specimen collected: LACM 70234, collected by J. BENEDICT 19 Jan. 1984; Northern Thailand, Aung Kang Region, Big House Cave; male, in alcohol with skull extracted. This single specimen was collected approximately one meter above a stream, on the cave wall. The field notes do not indicate whether or not this specimen was near other bats. This specimen represents the first record for this species outside of mainland China and extends the known range for this species by 1250 km southwest. The two other known localities are Omi San, Szechwan Province, China (the type locality) and 3 mi. east of Kweiyang City in Kweichow Province, China (STAGER 1949). Two of STAGER's six specimens (LACM 8214 and 8215) compare favorably with the new specimen. Selected measurements of LACM 70234, 8214, 8215, and the type (as given by THOMAS 1911) respectively are as follows: head and body length, 55, 60, 58, 55; forearm length, 44, 45.35, 42.90, 45; tibia, 16.50, 18.65, 18.10, 29 (reported as lower leg plus foot), hindfoot length, 11, 12, 12, none reported for type, ear length, 22, 24, 24, 22; greatest length of skull, 15.50, 15.95, 16, 15.2.

In addition to *M. altarium*, seven other species of *Myotis* are known from northern Thailand (LEKAGUL and MCNEELY 1977): *M. mystacinus*, *M. annectans*, *M. rosseti*, *M. siligorensis*, *M. horsfieldii*, *M. chinensis*, and *M. hasseltii*. Published information on the relationships of *M. altarium* are contradictory because so little material is available.

THOMAS (1911) stated that the morphology of *M. altarium* most resembled *M. pequininus*, a member of the large footed subgenus, *Leuconoe*. TATE (1941) placed *M. altarium* in the *M. emarginatus* section of the subgenus *Selysius*. The other species in this section of *Selysius* (sensu TATE 1941) are *M. emarginatus*, *M. peytoni* (= *montivagus*; HONACKI et al. 1982), *M. primula* (= *annectans*; HONACKI et al. 1982), and *M. saturatus* (= *emarginatus*; CORBET 1978). FINDLEY (1972) also placed *M. altarium* in the subgenus *Selysius*, but this analysis placed *M. altarium* close to a different group of species: *M. mystacinus*, *M. siligorensis*, *M. davidii* (= *mystacinus*; HONACKI et al. 1982), and *M. ikonnikovi*, FINDLEY's (1972) analysis and later classification placed the species *M. emarginatus* and *M. pequininus* into the subgenus *Myotis*. The relationships of *M. altarium* are

therefore confused, because according to TATE (1941) it should be placed close to *M. emarginatus* and so should be in the subgenus *Myotis* as this subgenus is characterized by FINDLEY (1972).

Based upon the three LACM specimens and THOMAS (1911), *M. altarium* can be characterized as follows: a medium sized *Myotis* with ears twice as long as they are wide, and which are distinctively black and translucent. The lateral edge of the ears are folded cranially. When laid forward the ears reach 5 mm beyond the tip of the muzzle. There is a distinct lobule at the base of the ear. The lobule is separated from the ear by a deep notch. The tragus is long, thin, and bluntly pointed. The flight membranes are naked, blackish brown in color and attach to the hind foot at the distal end of the metatarsal of digit one. There are eight striae on the uropatagium, which is naked dorsally, but with scattered hairs ventrally. The dorsal fur is light brownish in color and 12–15 mm in length. The ventral fur color is lighter having dark bases and whitish tips. The calcar is weakly lobed at its mid point. The hind foot measures over 60% of tibial length.

The skull of *Myotis altarium* (Fig. 1) has a short and distinctly upturned rostrum in lateral view. A line drawn along the long axis of the rostrum intersects a line drawn along the long axis of the cranium at approximately 145°. The skull slopes sharply anteriorly to a point above the posterior premolars. The skull profile has a slight convexity at mid-braincase (where the frontal and parietal bones meet). The nasal bones are flattened proximally and slightly arched (lateral to medial) distally and curved upwards distally at the premaxillary-nasal junction. The junction of the premaxillary-nasal bones is at the level of the superior surface of the orbit. The posterior part of the premaxillary bones are arched and display a obvious bump in lateral view. The zygomatic arches are consistant in depth, as seen in lateral view and expand somewhat posteriorly in dorsal view. The anteorbital foramen opens 0.7 mm anterior to the orbit above the anterior labial root of P³ and is 0.3 mm in diameter. The anteorbital foramen leads to an enclosed basin found above the roots of the premolars.




Fig. 1. Dorsal, lateral, and ventral views of the cranium of *M. altarium*. (Approximately 3× life size)

The ventral skull surface displays several distinctive features. The palate is deeply vaulted. An unusual feature of the palate is a sharp ridge running transversely between the canine teeth. The foramen ovale is large and conspicuous in ventral view. The basisphenoid is raised above the floor of the interbullar area resembling a bridge. There is no sagittal crest on any of the specimens. All upper and lower premolars are fully in line with the